

**Color Evolution in Moderate-Redshift Galaxy Clusters**

S.A. Stanford, IGPP/Livermore

P.R.M. Eisenhardt, JPL

M.E. Dickinson, STScI

We have obtained optical-IR imaging of  $\sim 40$  galaxy clusters in the range  $0.02 < z < 0.9$ . Our sample traces cluster galaxy properties over nearly half a Hubble time. The wide-field imaging typically reaches 2 mag below  $K_*$ , yielding  $\sim 50$ – $100$  galaxies per cluster. We are using this dataset in conjunction with archival WFPC2 imaging to investigate color evolution of the elliptical galaxy population. Analysis of a subset of the clusters has been carried out using IR-selected galaxy samples. After field-correction, we calculated median optical-IR colors of the morphologically-identified ellipticals. The colors are then zeropointed to the Coma cluster using our own  $UBVRIZJHK$  photometry of an IR-selected sample. The resulting observed-frame color differences represent change from the present epoch, largely independent of model-dependent  $k$ -corrections. We find a bluing trend with redshift in the optical-IR colors, in general agreement with the standard passive evolution scenario in which ellipticals are a coeval population formed at  $z > 5$ .

---

\*This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.